

Hyperspectral Remote Sensing of the Coastal Ocean: Adaptive Sampling and Forecasting of In situ Optical Properties

Mark Moline

Biological Sciences Department, California Polytechnic State University

San Luis Obispo, CA, 93407, U.S.A.

phone: (805) 756-2848 fax: (805) 756-1419 email: mmoline@calpoly.edu

Award #: N000149910197

<http://marine.rutgers.edu/cool/hycode2/hycode2.html>

LONG-TERM GOAL

We are developing an integrated rapid environmental assessment capability that will be used to feed an ocean nowcast/forecast system. The goal is to develop a capacity for predicting the dynamics in inherent optical properties in coastal waters. This is being accomplished by developing an integrated observation system that is being coupled to a data assimilative hydrodynamic bio-optical ecosystem model. The system was used adaptively to develop hyperspectral remote sensing techniques in optically complex nearshore coastal waters.

OBJECTIVES

Our objectives were to 1) develop and deploy moored, shipboard, and autonomous bio-optical systems in the coastal ocean to ground-truth remote sensing imagery, 2) use rapid environmental assessment techniques to quantify the physical, chemical and biological processes that define the spatial and temporal variability in the spectral IOPs for the nearshore coastal ocean during summer-time upwelling, 3) refine and calibrate existing hyperspectral optical models to derive IOPs from remotely sensed data using the above datasets and, 4) in collaboration with other principal investigators couple a radiative transfer ecosystem module to the data-assimilative hydrodynamic model. Our analysis on inverting signatures to bulk *in situ* IOP constituents, use derived phytoplankton signatures as inputs to bio-optical productivity models, define how the inherent optical properties impact both operational horizons of emergent light fluxes and remote sensing reflectance measured by the international constellation of satellites and aircraft. Finally we are conducting hindcast studies with coupled optical/physical numerical model.

APPROACH

We have conducted a series of Coastal Predictive Skill Experiments (CPSE) at the Long-term Ecosystem Observatory (LEO-15) in order to understand the physical forcing of the nearshore optical properties. To this end, coordinated shipboard (physical and bio-optical) and AUV adaptive sampling surveys of the upwelling centers were conducted based on the real-time remote observations and the model forecasts. The observational capability is used then to characterize and model the variability in the physics and the

associated in-water optical properties in order to ground-truth hyperspectral imagery collected by COIS sensors mounted on aircraft.

WORK COMPLETED

The field-work is accomplished at LEO-15, however we are pursuing similar efforts for calibration of aircraft and *in situ* hyperspectral sensors off California in October, 2002. The LEO-15 experiments were focused around dynamical forecasts generated through the Regional Ocean Model (ROMs) forecasts. These forecasts robustly predicted alternating upwelling/downwelling events. These forecasts, real-time CODAR fields, and *in situ* data from the autonomous nodes assisted in choosing flight missions for the aircraft (PHILLS 1, PHILLS 2, AVIRIS, Proteus, SPECTIR) and position three ships under the aircraft for *in situ* validation. Furthermore the *in situ* nodes were outfitted to measure a full suite of inherent optical properties. Shipes were similarly outfitted. Field sampling was coordinated through the modeling/observation system and allowed for 1) 16 clean overflights providing hyperspectral ocean color data with complete remote sensing ground truth data from the research fleet, 2) 5 days with more than two aircraft flying at one time allowing for unique vicarious calibration between aircraft systems, and 3) calibration of atmospheric parameters using NASA-funded aircraft. During the three years of field work in excess 700 discrete samples were collected and have been analyzed in the lab for filter pad absorption spectra, particle size, phytoplankton pigmentation, and organic carbon/nitrogen content.

The ROMS systems were coupled to the EcoSim model. This required substantial modification of the EcoSim model, which was not written in to run in parallel. ROMS is a free-surface, hydrostatic, primitive equation model initially based on the s-coordinate Rutgers University Model (SCRUM) described by (Song et al., 1994). ROMS was rewritten by the UCLA and Rutgers ocean modeling groups to improve its numerics and efficiency in single and multi-threaded computer architectures. New features include high-order advection schemes; accurate pressure gradient algorithms; several subgrid-scale parameterizations; atmospheric, oceanic, and benthic boundary layers; radiation boundary conditions; and data assimilation. EcoSim, which is upper ocean ecological model, was successfully coupled to the ROMS model and the shared memory parallel version (using OpenMP) of the ROMS/EcoSim code that we used to produce forecasts on the New Jersey coast for ONR's COMOP/HyCODE program (Arango et al., 2002) in July 2001 required 25 hours of wall clock time to process a 31 day simulation of a 100 x 240 (horizontal) x 25 (vertical) grid when executing on 16 (out of 256) processors of the NRL Origin3800 supercomputer (neo.cmf.nrl.navy.mil). Additionally, Ecolight, a version of the Hydrolight 4.1 radiative transfer model, was added to the ROMS/EcoSim modeling system. The advantage of adding Ecolight is that it provides an accurate solution of the RTE for any water body, given the absorption and scattering properties of the water body, the incident sky radiance, and the bottom reflectance (for finite-depth waters) but it runs one thousand times faster than the standard version of Hydrolight.

RESULTS

FIELD DATA During the experiment we experienced a wide gradient of hydrographic conditions, which provided a wide dynamic range ideal for calibrating the aircraft

hyperspectral sensors. The adaptive sampling strategy allowed us to optimize this calibration exercise, with over 10 days of high quality *in situ* and aircraft data. Cal Poly State University have been comparing the *in situ* and aircraft optical data collected by FERI. The conditions that dominated the optical signals varied between the two years. Summer 2000 was dominated by a large plume of Hudson River water. In contrast alternating upwelling and downwelling events dominated summer 2001.

Inversion of in situ data and optical tagging of water masses: We developed a method that takes ac-9 data and inverts the bulk signal into its constituent end members. The inversion method estimates the coefficients (w_i) and exponential slopes (s , r) using a non-linear constrained least-squares regression. Derived particulate, detrital and phytoplankton loads were compared to independent *in situ* and discrete data. Good agreement was achieved between measured and modeled particulate and phytoplankton spectra. Therefore the derived particulate spectra could be quantitatively derived from the ac-9 with no major bias as a function of wavelength. Comparing to HPLC data the inverter could predict ($p < 0.05$) the concentration of chlorophyll c and phycobilin-containing phytoplankton. We compared the water masses delineated optically from these derived products to those predicted with temperature and salinity in an effort to extending classical T-S water mass analysis into multi-dimensional space using derived optical parameters that ultimately could be derived using remote sensing platforms (aircraft or satellites). The T-S relationships detected the presence of significantly different water masses within and between each year, however their boundaries were difficult to resolve from T-S structure alone. While these optical parameters are semi-conservative, the relative change of these parameters is longer than many coastal ocean processes enhancing their potential utility. Examples of features that were easy to determine from HyCODE include the Hudson River plume with the high concentration of large particles, phytoplankton, CDOM and detritus during summer 2000. Here the river was dominated with shallow CDOM slopes consistent with terrestrially derived materials. A second example was dense bottom water, which was easily identified optically by steep CDOM slopes. The water mass clusters for the temperature/salinity and the optics (CDOM slope & load, detritus slope and load, and phytoplankton load) generally agreed with each delineating 3 major water masses in summer 2000 and 2 major water masses in 2001.

Bio-optical models for primary productivity. For summer 2000 we used the optical profiling node to collect time series. Using the ac-9 inversion method the spectral absorption of phytoplankton. The bulk absorption, attenuation and backscatter data from the optical profiler were used as input into a radiative transfer model (Hydrolight v. 4.2) to model the spectral scalar irradiance from 400-700 nm. These spectral were combined with the modeled phytoplankton absorption spectra from the OSI model to compute the spectrally-weighted absorption coefficient (\bar{a}_{ph}). The estimates of \bar{a}_{ph} were used as input to compute primary productivity. Modeled integrated water column productivity were within 11% of measured productivity results using ^{14}C incubations.

Effect of Phytoplankton Community Structure on inherent optical properties. High variability in physical, biological and chemical properties in coastal waters have limited our ability to sample the appropriate time and space scales to resolve physical forcing of

the ecosystem. To improve our understanding, the relationship between episodic summertime upwelling and downwelling events and the corresponding dynamics in bulk phytoplankton biomass and community structure was examined. Inherent and apparent optical properties were concurrently measured to evaluate the use of optics to improve future sampling coverage in coastal regions. Results indicate peak chlorophyll biomass tracked the maximum density gradient, and that increasing surface phytoplankton biomass was associated with decreasing stratification offshore over time. Diatoms dominated the study site, however significant shifts in cyanobacteria and dinoflagellate communities were observed. Dinoflagellate and cyanobacteria communities responded inversely to episodic events, with cyanobacteria being favored during intense downwelling. Differences in phytoplankton absorption properties significantly changed the corresponding in water inherent optical properties, allowing for characterization of the community structure from measurements of above water hyperspectral reflectance.




Figure 11

Figure 1. Results of the canonical correlation between the phytoplankton community abundance and the measured hyperspectral reflectance (R). The coefficients for the R wavelengths are shown for the (A) first, (B) second, (C) third and (D) fourth canonical correlates. The four correlates were significant ($p \ll 0.004$, $n = 91$) at 0.97, 0.93, 0.92 and 0.91 respectively. Significant R wavelengths were determined by a general linear model (see Methods) and are shown as $p < 0.01$ (white bars), $p = 0.01$ to 0.05 (gray bars) and $p > 0.05$ (black bars). Significant phytoplankton groups are also shown for each significant correlate as $p < 0.01$ (bold) and $p < 0.05$ (italics). For illustration, R measurements taken within 1 km of the shoreline on July 13th and August 1st, 2001 are overlaid in panels A and B, respectively.

IMPACT/APPLICATIONS

An integrated system for predicting the 3-dimensional structure of coastal currents, water density and in-water optical properties on the time scales of days is essential to numerous naval operations such as mine counter measures, special forces operations, amphibious landings, and shallow water anti-submarine warfare. The NEMO system is being designed to provide hyperspectral ocean color data for mapping in-water constituents in areas of high naval interest and the derived algorithms. Finally hydrodynamic/optical forecasting system provides the key to integrate and forecast the observed optical properties over time. HyCODE has played a central role to developing optical REMUS AUV and optical Webb Glider. All these observation and modeling systems are relocatable and will be key for future naval operations and homeland defense.

TRANSITIONS

The data is being freely shared. Data will be disseminated to the ONR WOOD database. Data is available via one-way FTP. The optical data is currently being utilized by NRL and NASA remote sensing projects. Finally the ongoing real-time data, for which the HyCODE program was central to for development, continues to be accessed via the web (over 65,000 hits/day) by the general public, Naval METOC groups, NAVO, NOAA Oil Spill response teams, and the U.S. Coast Guard.

RELATED PROJECTS

There were over 27 major institutional partners during the 2000-2001 experiments a large number supported by the HyCODE program. These efforts also complemented other independent efforts such as 1) validation of NAVAIR's KSS Lidar system, 2) ONR-YIP funded AUV bioluminescence prediction efforts, 3) ONR-STTR sponsored efforts to develop a "smart" fleet of automated Webb Gliders, 4) SeaSpace Inc. efforts to intercalibrate the international constellation of ocean color satellites, 5) calibration and refinement of a suite of NRL-derived satellite algorithms, 6) calibration of atmospheric parameters with NASA's atmospheric Chesapeake Lighthouse and Aircraft Measurements for Satellites experiment, 7) field infrastructure for NASA's NIP and PECASE remote sensing projects, and 8) model development for ONR's CBLAST Program.

PUBLICATIONS

(** papers by Cal Poly State University undergraduates and graduate students)

Schofield, O., Gryzmski, J., Bissett, P., Kirkpatrick, G., Millie, D. F., Moline, M. A.

Roesler, C. 1999. Optical monitoring and forecasting systems for harmful algal blooms: Possibility or pipedream? *Journal of Phycology*. 35: 125-145. [published, refereed]

- Kirkpatrick, G., Millie, D. F., Moline, M. A., Schofield, O. 2000. Absorption-based discrimination of phytoplankton species in naturally mixed populations. *Limnology and Oceanography* 42: 467-471. [published, refereed]
- Bissett, P., Schofield, O., Mobley, C. Crowley, M. F., Moline, M. A. 2001. Optical remote sensing techniques in biological oceanography. In: Methods in Marine Microbiology. Paul, J. (ed). Academic Press, London. 30: 519-538. [published, refereed]
- Moline, M. A., E. Heine**, J. Case, C. Herren and Schofield O. 2001. Spatial and temporal variability of bioluminescence potential in coastal regions. In: (J. F. Case, P. J. Herring, S. H. D. Haddock, L. J. Kricka and P. E. Stanley eds.) Bioluminescence and Chemiluminescence 2000. World Scientific Publishing Company, Singapore, pp. 123-126. [published, refereed]
- Grzymiski, J., **Orrico, C., Schofield, O. 2001. Monochromatic ultraviolet light induced damage to photosystem II efficiency and carbon fixation in the marine diatom *Thalassiosira pseudonanna* (3H). *Photosynthesis Research*. 68: 181-192. [published, refereed]
- **Oliver, M. 2001. Estimation of Phytoplankton production in dynamic coastal waters: An integrated physical, bio-optical and physiological approach. Masters Thesis. California Polytechnic State University. [published, refereed]
- Moline, O., Schofield, O., Grzymiski, J. 2002. Impact of dynamic light and nutrient environments on phytoplankton communities in the coastal ocean. In: Modeling Dynamic Systems: Dynamic Modeling for Marine Conservation Ecological Understanding. Lindholm, J. and Ruth, M. (eds) Springer Verlag 144-163. [published, refereed]
- Schofield, O., Bergmann, T., Bissett, W. P., Grassle, F., Haidvogel, D., Kohut, J., Moline, M., Glenn, S. 2002. Linking regional coastal observatories to provide the foundation for a national ocean observation network. *Journal of Oceanic Engineering*. 27(2): 146-154. [published, refereed]
- **Blackwell, S. M. 2002. A new AUV platform for studying bioluminescence in the coastal ocean. Masters Thesis. California Polytechnic State University. [published, refereed]
- **Heine, E. 2002. Mechanisms Structuring Bioluminescence in Near-Shore Waters Masters Thesis. California Polytechnic State University [published, refereed]
- **Orrico, C. 2002. *In situ* classification of phytoplankton in optically complex coastal waters. Masters Thesis. California Polytechnic State University. [published, refereed]
- Moline, M.A., Arnone, R., Bergmann, T., Glenn, S., **Oliver, M., **Orrico, C., Schofield, O., Tozzi, S. Variability in spectral backscatter estimated from satellites and its relation to in situ measurements in optically complex coastal waters. *Journal of International Remote Sensing*. 25(7-8): 1465-1468. [published, refereed]

- Tozzi, S., Schofield, O., Bergmann, T., Moline, M. A., Arnone, R. Variability in measured and modeled remote sensing reflectance for coastal waters at LEO-15. *Journal of International Remote Sensing*. 25(7-8): 1469-1472. [published, refereed]
- Chang G. C., T. D. Dickey, O. Schofield, A. D. Weidemann , E. Boss, M. A. Moline and S. M. Glenn. 2002. Nearshore physical forcing of bio-optical parameters in the New York Bight. *J. Geophysical Research*, ref. 10.1029/2001JC001018. [published, refereed]
- ** Blackwell, S., J. Case, S. Glenn, J. Kohut, M. A. Moline, M. Purcell, O. Schofield and C. VonAlt. 2002. A new AUV platform for studying near shore bioluminescence structure. In: *Proceedings of the 12th International Symposium on Bioluminescence and Chemiluminescence*, Herring, P. J., L. J. Kricka and P. E. Stanley, editors. World Scientific, London, pp. 197-200. [published, refereed]
- Chang, G., K. Mahoney, A. Briggs-Whitmire, D. D. R. Kohler, C. D. Mobley, M. Lewis, M. A. Moline, E. Boss, M. Kim, W. Philpot and T. D. Dickey. The new age of hyperspectral oceanography. *Oceanography* 17(2): 16-23. [published, refereed]
- Schofield, O., Arnone, R., Bissett, W. P., Dickey, T., Davis, Curt, Finkel, Z., Oliver, M. J., Moline, M. A. 2004. Watercolors in the coastal zone: What can we see? *Oceanography* 17(2): 24-43. [published, refereed]
- Oliver, M., J. Kohut, A. Irwin, O. Schofield. S. Glenn, M. A. Moline, and W.P. Bissett 2004. Bioinformatic approaches for objective detection of water masses. *Journal Geophysical Research – NEOS Special Section*. Vol 109 C07S04 doi: 10.1029/2003JC002072. [published, refereed]
- **Oliver, M. J., O. Schofield, T. Bergmann, S. Glenn, M. A. Moline and C. Orrico**. 2004. In-situ optically derived phytoplankton absorption properties in coastal waters and its utility for estimating primary productivity rates. : *Journal of Geophysical Research*. 109, C07S11, doi: 10.1029/2002JC001627. [published, refereed]
- Moline, M. A., S. Blackwell**, R. Chant, M. J. Oliver**, T. Bergmann, S. Glenn and O. M. E. Schofield. Episodic physical forcing and the structure of phytoplankton communities in the coastal waters of New Jersey. *Journal of Geophysical Research*. [in press]
- Glenn, S., R. Arnone, T. Bergman, P. Bissett, M. Crowley, J. Cullen, J. Gryzmski, D. Haidvogel, J. Kohut, M. Moline, M. Oliver**, C. Orrico**, R. Sherrell, T. Song, A. Weidemann, R. Chant, O. Schofield. The Biogeochemical Impact of Summertime Coastal Upwelling on the New Jersey Shelf. *Journal of Geophysical Research*. [in press]
- Schofield, O., T. Bergmann, W. P. Bissett, A. Irwin, M. Oliver**, C. Orrico**, M. A. Moline. Inversion of the Inherent Optical Properties and Their Utility for Delineation of Water Masses in Turbid Coastal Waters. *Journal of Geophysical Research*. [in press]

HONORS/AWARDS/PRIZES

M. A. Moline	Cal Poly State University	New Investigator Program Award	NASA
M. A. Moline	Cal Poly State University	Young Investigator Program Award	ONR
M. A. Moline	Cal Poly State University	Presidential Early Career Award for Scientists and Engineers (PECASE)	ONR